



From Wells to Burners: Methane Emissions from California Natural Gas

June 7st, 2016

Marc L. Fischer (mlfischer@lbl.gov)

◆ Problem Overview


- ❑ Significance of Natural Gas (NG) Methane
- ❑ Bottom-up Estimates of California NG methane emissions

◆ CALGEM-NG measurements

- ❑ Regional NG Emission Measurements for SF Bay Area
- ❑ UCD Airborne Measurements of NG Facilities
- ❑ LBNL Mobile Plume Integration (MPI) Measurements
- ❑ LBNL Residential Building and Appliance Measurements

◆ Summary and Recommendations

CALGEM team & collaborators



Imperial College
London



PICARRO JPL



LBNL: Seongeun Jeong, Toby Walpert, Xinguang Cui, Justin Bagley, Wanyu Chang, Woody Delp, Dev Millstein

BAAQMD: Abhinav Guha, Phil Martien, David Fairley, Saffet Tanrikulu

CARB: Ying-Kuang Hsu, Mathias Falk, Abhilash Vijayan, Jorn Herner, Bart Croes, Vernon Hughes, Larry Hunsaker, Marc Vayssières, Richard Bode, Anny Huang, Joseph Fischer, Jim Narady, Webster Tassat, Mac McDougall, Steve Rider, Steven Aston, Neil Adler, and Harlan Quan

CEC: Guido Franco, Yu Hou

CIT: Sally Newman, Paul Wennberg, Christian Frankenberg

EarthNetworks: Christopher D. Sloop

Kings College London: Heather Graven, Kieran Brophy

JPL: Nick Parazoo, Riley Duren, Chip Miller

LLNL: Tom Guilderson

NASA-Ames: Laura Iraci, Matthew Johnson, Emma Yates

NOAA-CCG: Arlyn Andrews, Laura Bianco, Ed Dlugokencky, Scott Lehman, John Miller, Jim Wilczak, Steve Montzka, Colm Sweeney, Pieter Tans

PG&E: François Rongere, Gerry Bong

Picarro: Chris Rella

Scripps/UCSD: Ralph Keeling, Jooil Kim, Ray F. Weiss

SJSU: Craig Clements, Neil Lareau, Matthew Lloyd

SNL: Ray Bamba, Hope Michelson, Brian LaFranci

UC Berkeley: Allen Goldstein

UCSB: Ira Leifer

UC Davis: Stephen Conley, Ian Faloona, Shobhit Mehrorta

UC Irvine: Don Blake, Xiaomei Xu

UCR: Jingsong Zhang

CALGEM
natural gas CH₄
research was
predominantly
supported by
the California
Energy
Commission
through grants
to LBNL and
UCD


Problem Overview

- Natural gas provides ~40% of California fossil fuel energy
- Methane is a potent short lived climate pollutant
 - 3% of well-to-burner NG leaked as methane approximately doubles climate forcing of remaining 97% gas combusted to CO₂ on 20 year timescale
- CA and US now moving to control CH₄ emissions
 - 10-20% of California's total methane emissions likely from NG
 - Entire production to consumption chain susceptible to emissions
 - Measurements now fill some gaps in understanding across NG infrastructure


Bottom-up Natural Gas CH₄ Emissions

- Map emissions w/ 2010/2011 US-EPA emission factors and CA specific GIS activity data
 - Production: Dry gas and petroleum wells
 - Transmission, compression, and storage
 - Distribution & consumption
- Estimated NG emissions ~ 330 Gg CH₄ yr⁻¹ (-20% to + 30% @ 95%)
 - Top-down studies in SoCAB suggest higher NG emissions (Peischl et al., 2013, Wunch et al., 2016)
 - 2016 US-EPA estimates increase production but decrease distribution emissions
 - NG is still small fraction of total California CH₄ emissions

CALGEM-NG Methane Flux (nmol m⁻² s⁻¹)




(Jeong et al., 2014)




CALGEM-NG CH₄ Measurements

- Regional Emissions
 - Tower measurements
 - Atmospheric Inversions
- Large Facilities
 - Aircraft Observations
- Localized Sources
 - Mobile Plume Integration
 - Building Studies

Collaborative tower measurements
Atmospheric Regional Inverse Modeling




LBNL Plume Integration



UC Davis Mass-balance UC Irvine VOC



LBNL Building Science



Regional NG Emission Estimate for San Francisco Bay Area


- Oct-Dec, 2015 sampling at six collaborative sites
 - CH₄, ethane, pentane, toluene, CO, and other VOC
 - Livermore hourly CH₄ & VOC
 - Daily flask sampling at other sites analyzed at NOAA and UCI
- Fossil VOC:CH₄ compositions adopted from PG&E gas reporting, airborne measurements, and previous mobile source studies (e.g., Kirchstetter et al., 1996)



SFBA Biological and Fossil CH₄ Sources

- Biological sources
 - Landfill 51%
 - Livestock 15%
 - Wastewater 6%
 - Wetland 3%
- Fossil sources
 - NG distribution 15%
(0.2% NG consumption)
 - Mobil and refining 4%

SF Bay CH₄ Emissions at 1 km



Sector Specific SFBay CH₄ Inversion


- Inversion of biological, NG, and petroleum CH₄ emissions constrained by CH₄ and VOCs
- Hierarchical Bayesian estimates optimize background offsets, VOC source compositions, and emission scaling factors

$$\begin{bmatrix} C_{CH_4} - C_{bg}^* \\ C_{C_2H_6} - C_{bg}^* \\ C_{iC_5H_{12}} - C_{bg}^* \\ C_{nC_5H_{12}} - C_{bg}^* \\ C_{C_7H_8} - C_{bg}^* \end{bmatrix} = \begin{bmatrix} FE_{S \setminus NG_{PL}} & FE_{NG} & FE_{PL} \\ 0 & FE_{NG} f_{C_2H_6_{NG}}^* & FE_{PL} f_{C_2H_6_{PL}}^* \\ 0 & 0 & FE_{PL} f_{iC_5H_{12}_{PL}}^* \\ 0 & 0 & FE_{PL} f_{nC_5H_{12}_{PL}}^* \\ 0 & 0 & FE_{PL} f_{C_7H_8_{PL}}^* \end{bmatrix} \begin{bmatrix} \lambda_{S \setminus NG_{PL}} \\ \lambda_{NG} \\ \lambda_{PL} \end{bmatrix}$$

Preliminary results:

- 1) Biological CH₄ dominates
 - 2) NG emissions higher than prior at 0.3-0.5% SFBA NG consumption (Jeong et al., in prep)
- Approach amenable to sustained observations in other locations

Posterior CH₄ by Sector




Facility Specific Emissions

San Joaquin Valley Production




- UCD Airborne mass balance measurements
- Example: April, 2014 Belridge South petroleum production field w/ steam injection
 - Clear downwind enhancements of CH_4 and ethane
- Emissions estimated from mass balance flight $1900 \pm 700 \text{ kg CH}_4 \text{ hr}^{-1}$ ($17 \pm 6 \text{ Gg CH}_4 \text{ yr}^{-1}$)
- Bottom-up $15 - 20 \text{ Gg CH}_4 \text{ yr}^{-1}$
- Collaborative observations of Kern River/Front fields show emissions varied with well completion (Leifer et al., in prep)



Facility Level Emissions: Natural Gas Storage

- UCD Airborne mass balance measurements
 - Four sites observed 3-8 times from June, 2014 – May, 2016 (+ four others recently)
 - Emissions vary from ND to $> 400 \text{ kg CH}_4 \text{ hr}^{-1}$
 - Median emissions $\sim 1 - 2 \times$ annual voluntary reporting
 - $\text{C}_2\text{H}_6:\text{CH}_4 \sim 5\%$ by vol., - consistent w/ NG
 - Single point failures carry high risk: Oct, 2015-Feb, 2016 Aliso Canyon well failure $\sim 30\%$ annual total CA fossil CH_4 emissions




(Mehrota et. al., in prep)

Facility Level Emissions: Petroleum Refining



- UCD Airborne mass balance measurements
 - Three refineries observed 3-5 times from Feb, 2015 – May, 2016
 - Emissions varied by site and date
~ 30 - 700 kg CH₄ hr⁻¹
 - Median emissions exceed annual emissions (4-25 x) reported to US-EPA
 - C₂H₆:CH₄ 6-10% by vol.



(Mehrota et. al., in prep)


Localized Source Emissions

LBL Mobile Plume Integration (MPI)

- Cross-wind integral of CH_4 enhancement flux quantifies localized plume emission
 - Sample inlets can be set to 4-8 m above ground
 - Multi-analyzer system w/ $^{13}\text{CH}_4$ allows NG attribution for strong plumes
 - Anemometry of wind velocity
 - Tests at LBNL and local utilities show 30% accuracy with 3 passes with steady winds & small obstructions



4 m
2 m
1 m




Localized Plume Measurements: Bakersfield Distribution & Consumption

- Survey 80km of Bakersfield public streets
- Detect 20 large (~ 1 ppm) leaks above background
- 40% of total emissions found within 0.5 km of large distribution pipes
- Plume integrations yield total emissions of $6.4 \text{ kg CH}_4 \text{ hr}^{-1}$
- Scaling by area suggests total emissions ~ $90 \text{ kg CH}_4 \text{ hr}^{-1}$
- Comparing with consumption suggests ~ 0.3% distribution leakage – similar to bottom up



CH₄ enhancements (green), distribution (orange) and transmission (blue) pipelines




Localized Plumes: Sacramento Delta Gas Wells


- CA Dept. Cons. well map data
- Initial inspection of 13 capped or idle wells
 - Quantify one plume $5 \pm 1.7 \text{ g CH}_4 \text{ hr}^{-1}$ (5 passes)
 - Detected three plumes $1.6\text{-}14 \text{ g CH}_4 \text{ hr}^{-1}$ (1pass each)
 - Non-detect downwind at 2 sites
 - 7 sites did not allow downwind access



Methane Plume



CH₄ Enhancement in vertical plane




Capped well

Localized Sources:


East Bay Distribution & Consumption

- Small ($\ll 1$ ppm) CH_4 plumes observed in SF East Bay
- Individual plumes emit $\sim 0.07 - 0.3 \text{ g CH}_4 \text{ hr}^{-1}$
- Emissions largest on commercial avenues w/ food service
- Total emissions $\sim 5 \text{ g CH}_4 \text{ hr}^{-1}$ over 30km route



Whole Building Measurements: Quiescent Residential Emissions


- Measurements
 - Depressurize house producing controlled inflow of outdoor air
 - Measure CH_4 enhancement relative to outdoor air
 - $^{13}\text{CH}_4/^{12}\text{CH}_4$ identifies NG vs. biological
- Results from 10 SF Bay homes
 - Median leak rate $0.2 \text{ g CH}_4 \text{ hr}^{-1}$ (0.1-0.4 $\text{g CH}_4 \text{ hr}^{-1}$ lower-upper quartiles)
 - Equivalent to $\sim 0.2\%$ of house consumption
- CEC project underway to measure 50-75 homes across CA housing stock



$$C_0 Q + E = C_i Q$$


$$E = Q (C_i - C_0)$$

Measured indoor (white) and outdoor (grey) methane during calibrated indoor leak (red)



Combustion Appliance Emissions

- Emission Ratio Method
 - Emission = product of $\text{CH}_4:\text{CO}_2$ enhancements * measured NG usage
- Tank-less water heaters
 - Test of three tank-less water heaters yield emissions of 3 - 12 $\text{g CH}_4 \text{ hr}^{-1}$
(1 hr operation ~ equal 1 day of quiescent house leakage)
- Clothes Driers and Gas Cooktop
 - One gas range emitted ~ 2 $\text{g CH}_4 \text{ hr}^{-1}$ in continuous operation
 - Two clothes driers emitted ~ 0.4 $\text{g CH}_4 \text{ hr}^{-1}$ emissions in continuous operation



Summary

1. CH₄ emissions present across all NG subsectors from wells to burners
2. Regional inversions suggest emissions from SFBA distribution ~ 0.3-0.5% of NG consumption
3. Production field measurements (limited but) ~ consistent with bottom-up but expect variability (particularly well completion)
4. Gas storage facility emissions variable but ~ consistent with reporting
5. Petroleum refining emissions appear larger (4-25 x) than reporting
6. Localized emissions in distribution & consumption sectors measurable and appear to ~ scale with gas throughput

Recommendations

1. Daily multi-species tower measurements needed for inversion-base verification of regional integrated NG CH₄ emissions
2. Plume imaging from ground, air, and space needed to identify local emission hotspots to guide site specific quantification and mitigation
3. Mass balance flights and mobile plume integration needed for quantitative assessment of facility and localized source emissions
4. Continuous (open-path or multi-point CH₄) sensing valuable for ongoing leak detection at high volume/flow facilities
5. Energy efficiency programs would benefit from added leak detection and repair procedures and revised standards guidance for low-emission appliances